

Geographic Variation in the Prevalence of Attention-Deficit/Hyperactivity Disorder: The Sunny Perspective

Martijn Arns, Kristiaan B. van der Heijden, L. Eugene Arnold, and J. Leon Kenemans

Background: Attention-deficit/hyperactivity disorder (ADHD) is the most common psychiatric disorder of childhood, with average worldwide prevalence of 5.3%, varying by region.

Methods: We assessed the relationship between the prevalence of ADHD and solar intensity (SI) (kilowatt hours/square meters/day) on the basis of multinational and cross-state studies. Prevalence data for the U.S. were based on self-report of professional diagnoses; prevalence data for the other countries were based on diagnostic assessment. The SI data were obtained from national institutes.

Results: In three datasets (across 49 U.S. states for 2003 and 2007, and across 9 non-U.S. countries) a relationship between SI and the prevalence of ADHD was found, explaining 34%–57% of the variance in ADHD prevalence, with high SI having an apparent preventative effect. Controlling for low birth weight, infant mortality, average income (socioeconomic status), latitude, and other relevant factors did not change these findings. Furthermore, these findings were specific to ADHD, not found for the prevalence of autism spectrum disorders or major depressive disorder.

Conclusions: In this study we found a lower prevalence of ADHD in areas with high SI for both U.S. and non-U.S. data. This association has not been reported before in the literature. The preventative effect of high SI might be related to an improvement of circadian clock disturbances, which have recently been associated with ADHD. These findings likely apply to a substantial subgroup of ADHD patients and have major implications in our understanding of the etiology and possibly prevention of ADHD by medical professionals, schools, parents, and manufacturers of mobile devices.

Key Words: ADHD, chronobiological, circadian, light, prevalence, solar intensity

Attention-deficit/hyperactivity disorder (ADHD) is the most common psychiatric disorder of childhood, with average worldwide prevalence of 5.3% (1). Heritability of ADHD has been reported as 60%–70% (2). Pre- and perinatal factors (preterm birth, low birth weight [LBW]) also play an important role in the etiology of ADHD (3). Additional factors might include global living conditions that could be investigated by inspecting geographic differences in ADHD prevalence.

Previous research reported that 78% of adult unmedicated (4) and one third of medication-naïve pediatric (5) ADHD patients have idiopathic sleep onset insomnia (SOI), accompanied by delayed circadian phase as measured by delayed dim light melatonin onset (6). This finding is supported by melatonin signaling deficiencies (7), clock gene abnormalities (8), and a higher prevalence of “evening types” in adult ADHD (9). Furthermore, a recent meta-analysis incorporating data from 35,936 healthy children reported that sleep duration correlates positively with school performance, executive function, and negatively with internalizing and externalizing behavior problems (10). In addition,

a sleep restriction regime for 3–7 days results in cumulative impairment of attention in healthy adults (11,12) and children (13) and in a tendency for increased θ electroencephalogram (EEG) power (effect size $d = .53$), (14), which is a measure reported in some ADHD studies, also reflective of fatigue or drowsiness (15,16). These impairments of attention did not normalize after a single night of recovery sleep but only after at least the same number of recovery nights as the number of sleep-restricted nights (11,12). Chronobiological interventions in ADHD patients with insomnia, such as early-morning bright light (17) and long-term melatonin treatment (18), have resulted in an improvement of ADHD symptoms. Additionally, improvement of ADHD symptoms was shown after treatment of other sleep disorders, such as dopamine agonist (19) or iron supplementation (20) for restless legs syndrome and adenotonsillectomy in sleep apnea by (21). Together, these results suggest that, at least in a subgroup of ADHD, the symptoms are exacerbated by sleep problems. In the aforementioned SOI studies, patients were unmedicated (4) or medication-naïve (5). Many well-controlled studies in unmedicated ADHD patients have reported these sleep problems [for review see Yoon *et al.* (22)]. Thus the increased prevalence of sleep problems in ADHD cannot be explained by effects of stimulant medication. Nevertheless, stimulants often aggravate sleep problems that already existed before stimulant use.

The association of ADHD with circadian disturbances and the powerful impact of bright light on the circadian rhythm, sleep, and daytime function led us to investigate in more detail the relationship between environmental light exposure and ADHD prevalence. This was further encouraged by a graph from the U.S. Centers for Disease Control and Prevention (CDC) (23) depicting differences in ADHD prevalence for the U.S. (Figure 1, left) and solar intensity (SI) maps for the U.S., available from the National Renewable Energy Laboratory (NREL) (Figure 1, right). We noted that the U.S. states with maximum SI (Arizona, New Mexico,

From the Department of Experimental Psychology (MA, JLK), Utrecht University, Utrecht; Research Institute Brainclinics (MA), Nijmegen; Leiden Institute for Brain and Cognition (KBvdH), Department of Clinical Child and Adolescent Studies, Leiden University, Leiden, the Netherlands; and Ohio State University (LEA), Columbus, Ohio.

Address correspondence to Martijn Arns, Ph.D., Department of Experimental Psychology, Research Institute Brainclinics, Bijleveldsingel 34, Nijmegen 6524 AD, The Netherlands; E-mail: martijn@brainclinics.com.

Received Dec 6, 2012; revised and accepted Feb 19, 2013.




Figure 1. Comparison between the prevalence of attention-deficit/hyperactivity disorder (ADHD) and solar intensity (SI) in the U.S. This figure demonstrates on the left map the prevalence rate of ADHD among different U.S. states (U.S. Centers for Disease Control and Prevention, 2010; <http://www.cdc.gov/ncbddd/adhd/prevalence.html>). The figure on the right depicts the horizontal irradiation or SI (SI in kilowatt hours/square meters/day) across the U.S. as reported by the National Renewable Energy Laboratory (map created by the National Renewable Energy Laboratory for the Department of Energy). Note the similarity between the areas with highest SI (depicted in red) and those with the lowest prevalence of ADHD (palest blue).

Nevada, California, Utah, and Colorado) also had the lowest prevalence of ADHD. Therefore, in this study we sought to systematically investigate the relationship between the prevalence of ADHD and SI as a measure of visible and nonimage forming light (expressed in kilowatt hours/square meters/day). We hypothesized that high SI is associated with lower rates of ADHD by its overwhelming phase-advancing effect on the circadian clock.

Methods and Materials

ADHD Prevalence Estimates

Searches were performed to uncover statistics about the prevalence (PREV) of ADHD around the world or across states employing standardized procedures, to facilitate valid comparisons of PREV estimates.

Solar Intensity

Horizontal irradiation (radiation reaching the surface of the earth on a horizontal plane, expressed in kilowatt hours/square meters/day; SI data: SI) were obtained from national and international agencies: U.S. National Renewable Energy Laboratory (NREL^A) for the U.S. (NREL: P. Gray-Hann, personal communication, August 24, 2012), global horizontal irradiation, State University of New York (Mexico), global horizontal irradiation, National Institute for Spatial Research (Colombia), and the European Institute for Energy and Transport (IET^B) for Europe and other countries.

The radiation model used by NREL was developed by Perez *et al.* (24) and "...uses hourly radiance images from geostationary weather satellites, daily snow cover data, and monthly averages of atmospheric water vapor, trace gases, and the amount of aerosols in the atmosphere to calculate the hourly total insolation...falling on a horizontal surface..." (25). Similar methods were used by IET; for more details see Šúri *et al.* (26).

Statistics

The relationship between PREV and SI was investigated by curve estimation. For nonlinear relationships, an appropriate

transformation (log or square root) was applied to obtain the most significant linear trend. A correlation was calculated between PREV and SI as well as with other potential confounding variables, such as latitude, altitude, and (for the U.S.) average income (socioeconomic status), LBW, and infant mortality (IM)/state (CDC data) and other factors reported by the ADHD PREV studies obtained (Pearson correlation, two-tailed). Those factors that were found significantly correlated with PREV were introduced as control variables in partial correlation analyses between PREV and SI (two-tailed). Finally, to investigate specificity for ADHD, we conducted the same analyses for autism spectrum disorders (ASD) and major depressive disorder (MDD) in the U.S. data that were also obtained from the CDC.

Results

Two data sources were identified that employed identical methods to estimate the prevalence of ADHD. These sources were the PREV estimates in children from the CDC/U.S. state (23) and the PREV estimates for adults across several countries (27).

ADHD Prevalence Across U.S. States: CDC

The CDC data were collected in 2003 and 2007 and were part of the National Survey of Children's Health in children under 18 years of age. This survey was a national, cross-sectional, random-digit-dialed landline telephone survey, where one child was selected randomly from each household to be the focus of the parent or guardian interview (response rates: 68.8% in 2003 and 46.7% in 2007) (23). Parents were asked whether or not a doctor or other healthcare provider had ever told them that their child had "attention-deficit disorder or attention-deficit/hyperactivity disorder, that is, ADD or ADHD" (23). Higher rates of ADHD were found among boys, multiracial children, and children covered by Medicaid.

Figure 2 demonstrates the relationship between SI and PREV for 2003 and 2007. As can be seen from Figure 2, this relationship does not seem linear, which was confirmed by curve estimation (all statistics are represented for 2003 | 2007, respectively). The best fit was obtained by a sigmoidal dose-response relation




Figure 2. The association between attention-deficit/hyperactivity disorder (ADHD) prevalence rates and solar intensity for different U.S. states for 2003 (left) and 2007 (right). A sigmoidal dose-response relation best describes the association. CDC, U.S. Center for Disease Control and Prevention; NREL, National Renewable Energy Laboratory.

[$F_{1,46} = 13.78$; $p = .0006$; $R^2 = .36$ | $F_{1,46} = 16.38$; $p = .0002$; $R^2 = .37$], and log transformation of PREV and SI resulted in the best linear fit with a significant correlation between PREV and SI [$p = .002$; $r = -.429$ | $p = .004$; $r = -.409$; $df = 49$]. The PREV (log transformed) did not correlate with socioeconomic status (all p values $< .199$) but did correlate with latitude for 2003 [$p = .050$; $r = -.281$; $df = 49$] but not for 2007 [$p = .094$; $r = -.242$; $df = 49$]. Furthermore, PREV was positively associated with percentage LBW [$p = .001$; $r = .468$; $df = 49$ | $p = .003$; $r = .422$; $df = 49$] and IM [$p = .001$; $r = .468$; $df = 49$ | $p = .001$; $r = .457$; $df = 49$], in line with the literature (3). Partial correlations including the aforementioned variables as covariates made the relationship between PREV and SI stronger [$p < .000$; $r = -.637$; $df = 44$ | $p < .000$; $r = -.580$; $df = 44$], resulting in explained variance of 34%–41% of ADHD prevalence.

To control for the demographic differences in ADHD PREV as reported by CDC (23), we also conducted partial correlations adding Medicaid coverage (CDC data), male/female ratio (census data) and percentage multi-raciality (census data)/state, which did not change the relationship between ADHD PREV and SI for 2003 or 2007 (all p values $\leq .003$). To control further for racial and ethnic differences, we also included the percentage Hispanic or Latino population/state (census data), which slightly lowered the correlation between SI and PREV but still resulted in a significant association [$r = -.443$; $p = .002$; $df = 43$ | $r = -.353$; $p = .017$; $df = 43$].

To test whether the aforementioned findings were moderated by seasonal fluctuations in light intensity, the aforementioned analyses were repeated with SI data/month and PREV (2003 or 2007). Partial correlations correcting for latitude, LBW, and IM resulted in significant correlations between all monthly SI values and ADHD PREV for 2003 and 2007 (all $p < .004$; $-.415 < r < -.661$), suggesting that differences in monthly SI did not influence the association SI with PREV.

When we repeated the aforementioned analysis excluding states with an SI > 5.5 (Arizona, New Mexico, Nevada, California, Utah, and Colorado, representing 22% of the total U.S. population on the basis of the 2010 U.S. census data) from the analysis, the partial correlation between SI and PREV weakened but remained significant for 2003 [$p = .041$; $r = -.324$; $df = 38$] and marginally significant for 2007 [$p = .082$; $r = -.278$; $df = 38$], suggesting these effects are mostly driven by the high SI in these six states. These six states are all characterized by a higher

elevation, so we checked for effect of high-altitude thin air. There was a significant correlation for the whole sample between PREV and elevation [$p < .000$; $r = -.672$; $df = 49$ | $p < .000$; $r = -.571$; $df = 49$] and between SI and elevation [$p < .000$; $r = .644$; $df = 49$]. However, partial correlations between PREV and elevation were no longer significant when adding SI as a covariate ($p < .4$), whereas adding elevation as a covariate still allowed marginally significant and significant associations between PREV and SI for 2003 and 2007, respectively ($p = .076$ | $p = .007$).

Prevalence rates were also available for ASD (14 U.S. States, 2008) and MDD (44 U.S. states, 2006–2008) from the CDC. The aforementioned analyses were repeated for ASD and MDD as well and did not result in any significant associations between PREV and SI ($p < .913$).

Non-U.S. Adult ADHD Prevalence Rates, Cross-National Data

In a multinational study by Fayyad *et al.* (27) identical methods were used to estimate the PREV of adult ADHD. A total of 11,422 respondents (18–44 years of age) from multi-stage household probability samples were screened by face-to-face surveys. Subjects were retrospectively assessed for childhood ADHD with the Diagnostic Interview Schedule for DSM-IV (28), and if subjects met childhood ADHD criteria, they were asked about whether they continued to have current problems with attention or hyperactivity-impulsivity. The response rate was 67.9% (27).

These authors reported a significantly lower PREV for Spain, Lebanon, Colombia, and Mexico and a significantly higher PREV for France as compared with Italy, Germany, Belgium, The Netherlands, and the U.S. sociodemographic effects on ADHD PREV were only found for a higher prevalence in men and among people educated less than at University level, albeit with modest magnitude ($1.5 < \text{odds ratio} < 3.0$), and these factors did not differ between countries. The four countries with lower prevalence also have the highest SI compared with the other countries, as can be seen in Figure 3. In this study we excluded the results from the U.S., because those have been analyzed in the preceding text in more detail.

A sigmoid dose-response was observed for the relation between PREV and SI [$F_{1,6} = 15.83$; $p = .0073$; $R^2 = .85$]. A log transformation of PREV and SI resulted in a linear relationship and a significant correlation between PREV and SI [$p = .018$;

This hypothesis is in line with recently reported results of morning bright light as a treatment for adults with ADHD (17) as well as with the sigmoidal dose-response effects for SI and PREV, which is also reported for circadian measures such as multi-unit activity of the suprachiasmatic nuclei in response to light intensity (37) and melatonin suppression as a result of exposure to light (38). Interestingly, in the CDC data the most significant increase in ADHD PREV between 2003 and 2007 was noted for adolescents 15–17 years old ($p < .001$) compared with 4–10-year ($p = .013$) and 11–14-year ($p = .016$) age groups. Furthermore, a recent study on the use of stimulant medication in the U.S. also found that across the last 12 years there was a slow but steady increase in pediatric stimulant use, primarily as a result of increased stimulant use in adolescents (39). We would expect that adolescents more often engage in modern media use in the evening as compared with younger children.

Controlled studies are required to prospectively replicate these findings and investigate whether intense light/solar exposure during the day, particularly in the morning, or reduced light exposure in the evening (especially blue 464–484-nm light) could reduce ADHD symptoms or maybe “treat” ADHD. This could open the way to prevention of a subgroup of ADHD or children with attention-deficit syndrome in a variety of ways: for example, exposing children more to natural light during the day (skylight to bring natural light into classrooms and scheduling more outside play time in the morning rather than in the afternoon); reducing exposure to blue light in the evening by parental control; or encouraging device manufacturers to control the emission of blue light from mobile devices on the basis of time of day. Furthermore, these results suggest that future genetic studies of ADHD might include genes involved in the circadian system and incorporate sleep data (e.g., actigraphy) and circadian parameters (dim light melatonin onset). These might define intermediate endophenotypes in the etiology of ADHD subgroups from gene \times environment interactions.

We acknowledge the National Renewable Energy Laboratory (NREL) and Pamela Gray-Hann for providing additional data on the U.S. solar potential/state and Marijtje Jongma, Ph.D., for her advice and support in the curve fitting procedures. Solar intensity data used in this article from NREL are available at <http://www.nrel.gov/gis/solar.html> (data accessed September 6, 2012) and from the European Institute for Energy and Transport are available at <http://re.jrc.ec.europa.eu/pvgis/apps/pvreg.php?lang=en&map=europe> (data accessed September 6, 2012).

MA initiated the article and has been involved with the data collection, data analysis, and manuscript writing and had full access to the data and takes responsibility for the integrity of the data and the accuracy of the data analysis. KVDH and JLK have contributed with suggestions for data analysis, interpretation of results, and manuscript editing at all stages. LEA has critiqued the ideas from the beginning, contributed several alternative hypotheses to be tested (e.g., elevation and vitamin D), and reviewed/edited several drafts of the article.

LEA has received research funding (to the university) or advisory board honoraria from AstraZeneca, Biomarin, CureMark, Lilly, Novartis, Noven, Seaside Therapeutics, and Shire and travel support from Noven.

MA, KBH, and JLK report no biomedical financial interests or potential conflicts of interest.

- Polaczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA (2007): The worldwide prevalence of ADHD: A systematic review and meta-regression analysis. *Am J Psychiatry* 164:942–948.
- Cortese S, Faraone SV, Sergeant J (2011): Misunderstandings of the genetics and neurobiology of ADHD: Moving beyond anachronisms. *Am J Med Genet B Neuropsychiatr Genet* 156:513–516.
- Halmøy A, Klungsoyr K, Skjærven R, Haavik J (2012): Pre- and perinatal risk factors in adults with attention-deficit/hyperactivity disorder. *Biol Psychiatry* 71:474–481.
- Van Veen MM, Kooij JJS, Boonstra AM, Gordijn MCM, Van Someren EJW (2010): Delayed circadian rhythm in adults with attention-deficit/hyperactivity disorder and chronic sleep-onset insomnia. *Biol Psychiatry* 67:1091–1096.
- Corkum P, Tannock R, Moldofsky H (1998): Sleep disturbances in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 37:637–646.
- Van der Heijden KB, Smits MG, Van Someren EJW, Gunning WB (2005): Idiopathic chronic sleep onset insomnia in attention-deficit/hyperactivity disorder: A circadian rhythm sleep disorder. *Chronobiol Int* 22: 559–570.
- Chaste P, Clement N, Botros HG, Guillaume J-L, Konyukh M, Pagan C, *et al.* (2011): Genetic variations of the melatonin pathway in patients with attention-deficit and hyperactivity disorders. *J Pineal Res* 51: 394–399.
- Baird AL, Coogan AN, Siddiqui A, Donev RM, Thome J (2012): Adult attention-deficit hyperactivity disorder is associated with alterations in circadian rhythms at the behavioural, endocrine and molecular levels. *Mol Psychiatry* 17:988–995.
- Bijlenga D, van der Heijden KB, Breuk M, van Someren EJW, Lie MEH, Boonstra AM, *et al.* (2011): Associations between sleep characteristics, seasonal depressive symptoms, lifestyle, and ADHD symptoms in adults [published online ahead of print December 29]. *J Atten Disord*.
- Astill RG, Van der Heijden KB, Van Ijzendoorn MH, Van Someren EJ (2012): Sleep, cognition, and behavioral problems in school-age children: A century of research meta-analyzed. *Psychol Bull* 138: 1109–1138.
- Axelsson J, Kecklund G, Åkerstedt T, Donofrio P, Lekander M, Ingre M (2008): Sleepiness and performance in response to repeated sleep restriction and subsequent recovery during semi-laboratory conditions. *Chronobiol Int* 25:297–308.
- Belenky G, Wesensten NJ, Thorne DR, Thomas ML, Sing HC, Redmond DP, *et al.* (2003): Patterns of performance degradation and restoration during sleep restriction and subsequent recovery: A sleep dose-response study. *J Sleep Res* 12:1–12.
- Fallone G, Acebo C, Seifer R, Carskadon MA (2005): Experimental restriction of sleep opportunity in children: Effects on teacher ratings. *Sleep* 28:1561–1567.
- Beebe DW, Rose D, Amin R (2010): Attention, learning, and arousal of experimentally sleep-restricted adolescents in a simulated classroom. *J Adolesc Health* 47:523–525.
- Arns M, Kenemans L (2012): Neurofeedback in ADHD and insomnia: Vigilance stabilization through sleep spindles and circadian networks [published online ahead of print October 23]. *Neurosci Biobehav Rev*.
- Arns M, Conners K, Kraemer H (2012): A decade of EEG theta/beta ratio research in ADHD: A meta-analysis [published online ahead of print October 19]. *J Atten Disord*.
- Rybak YE, McNeely HE, Mackenzie BE, Jain UR, Levitan RD (2006): An open trial of light therapy in adult attention-deficit/hyperactivity disorder. *J Clin Psychiatry* 67:1527–1535.
- Hoebert M, van der Heijden KB, van Geijlswijk IM, Smits MG (2009): Long-term follow-up of melatonin treatment in children with ADHD and chronic sleep onset insomnia. *J Pineal Res* 47:1–7.
- Cortese S, Lecendreux M, Bernardina BD, Mouren MC, Sbarbati A, Konofal E (2008): Attention-deficit/hyperactivity disorder, Tourette's syndrome, and restless legs syndrome: The iron hypothesis. *Med Hypotheses* 70:1128–1132.
- Walters AS, Mandelbaum DE, Lewin DS, Kugler S, England SJ, Miller M (2000): Dopaminergic therapy in children with restless legs/periodic limb movements in sleep and ADHD. Dopaminergic therapy study group. *Pediatr Neurol* 22:182–186.

21. Huang Y-S, Guilleminault C, Li H-Y, Yang C-M, Wu Y-Y, Chen N-H (2007): Attention-deficit/hyperactivity disorder with obstructive sleep apnea: A treatment outcome study. *Sleep Med* 8:18–30.
22. Yoon SY, Jain U, Shapiro C (2012): Sleep in attention-deficit/hyperactivity disorder in children and adults: Past, present, and future. *Sleep Med Rev* 16:371–388.
23. Visser SN, Bitsko RH, Danielson ML, Perou R, Blumberg SJ (2010): Increasing prevalence of parent-reported attention-deficit/hyperactivity disorder among children—United States, 2003 and 2007. *MMWR Morb Mortal Wkly Rep* 59:1439–1443.
24. Perez R, Ineichen P, Moore K, Kmiecik M, Chain C, George R, Vignola F (2002): A new operational model for satellite-derived irradiances: Description and validation. *Solar Energy* 73:307–317.
25. National Renewable Energy Laboratory (2012): *Solar Maps Development—How the Maps Were Made*. Available at: http://www.nrel.gov/gis/solar_map_development.html. Accessed November 27, 2012.
26. Šúri M, Huld TA, Dunlop ED, Ossenbrink HA (2007): Potential of solar electricity generation in the European Union member states and candidate countries. *Solar Energy* 81:1295–1305.
27. Fayyad J, De Graaf R, Kessler R, Alonso J, Angermeyer M, Demyttenaere K, *et al.* (2007): Cross-national prevalence and correlates of adult attention-deficit hyperactivity disorder. *Br J Psychiatry* 190: 402–409.
28. Robins LN, Cottler LB, Buchholz KK, Compton W (1995): Diagnostic interview schedule for DSM-IV (DIS-IV). St. Louis, Missouri: Washington University.
29. Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, *et al.* (2003): The epidemiology of major depressive disorder: Results from the national comorbidity survey replication (NCS-R). *JAMA* 289: 3095–3105.
30. Even C, Schröder CM, Friedman S, Rouillon F (2008): Efficacy of light therapy in nonseasonal depression: A systematic review. *J Affect Disord* 108:11–23.
31. Getahun D, Jacobsen SJ, Fassett MJ, Chen W, Demissie K, Rhoads GG (2013): Recent trends in childhood attention-deficit/hyperactivity disorder. *JAMA Pediatr* 21:1–7.
32. Tolppanen A-M, Sayers A, Fraser WD, Lewis G, Zammit S, Lawlor DA (2012): The association of 25-hydroxyvitamin d(3) and d(2) with behavioural problems in childhood. *PLoS One* 7:e40097.
33. Custers K, Van den Bulck J (2012): Television viewing, internet use, and self-reported bedtime and rise time in adults: Implications for sleep hygiene recommendations from an exploratory cross-sectional study. *Behav Sleep Med* 10:96–105.
34. Van den Bulck J (2004): Television viewing, computer game playing, and internet use and self-reported time to bed and time out of bed in secondary-school children. *Sleep* 27:101–104.
35. Wood B, Rea MS, Plitnick B, Figueiro MG (2013): Light level and duration of exposure determine the impact of self-luminous tablets on melatonin suppression. *Appl Ergon* 44:237–240.
36. Cajochen C, Frey S, Anders D, Späti J, Bues M, Pross A, *et al.* (2011): Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *J Appl Physiol* 110:1432–1438.
37. Meijer JH, Watanabe K, Schaap J, Albus H, Détári L (1998): Light responsiveness of the suprachiasmatic nucleus: Long-term multiunit and single-unit recordings in freely moving rats. *J Neurosci* 18:9078–9087.
38. Brainard GC, Hanifin JP, Greenson JM, Byrne B, Glickman G, Gerner E, Rollag MD (2001): Action spectrum for melatonin regulation in humans: Evidence for a novel circadian photoreceptor. *J Neurosci* 21:6405–6412.
39. Zuvekas SH, Vitiello B (2012): Stimulant medication use in children: A 12-year perspective. *Am J Psychiatry* 169:160.